

Datasheet: MCA1908HT

Description:	MOUSE ANTI HUMAN EPITHELIAL MEMBRANE ANTIGEN
Specificity:	EPITHELIAL MEMBRANE ANTIGEN
Other names:	CD227, MUCIN 1
Format:	Ready To Use
Product Type:	Monoclonal Antibody
Clone:	E29
Isotype:	IgG2a
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	■			Neat
Immunohistology - Paraffin	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Tissue Culture Supernatant - liquid , ready to use
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Immunogen	Delipidated extract of human cream
External Database Links	<p>UniProt: P15941 MUC1_HUMAN</p> <p>Entrez Gene: 4582 MUC1</p>
Synonyms	PUM
Specificity	MCA1908HT recognises epithelial membrane antigen (EMA), a heavily glycosylated protein of 265-400kD. This antibody stains normal and neoplastic epithelium including adenocarcinomas from a range of sites, squamous and transitional cell carcinomas and mesotheliomas. Strong staining of glandular epithelium is seen, but staining of squamous epithelium may be rather patchy. Positive staining with this antibody demonstrates epithelial origin of tumours.
Immunohistology	This product does not require protein digestion pre-treatment of paraffin sections. This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections.

Histology Positive Control Tissue Kidney

References

1. Cordell, J. *et al.* (1985) Production of monoclonal antibodies against human epithelial membrane antigen for use in diagnostic immunocytochemistry. [Br. J. Cancer. 52: 347-54.](#)
2. Gusev, Y. *et al.* (2001) Rolling circle amplification: a new approach to increase sensitivity for immunohistochemistry and flow cytometry. [Am. J. Pathol. 159 \(1\): 63 - 69.](#)
3. ten Berge, R.L. (2001) MUC1 (EMA) is preferentially expressed by ALK positive anaplastic large cell lymphoma, in the normally glycosylated or only partly hypoglycosylated form. [J Clin Pathol. 54: 933-9.](#)
4. Cao, Y. *et al.* (1998) Immunohistochemical characterization of a panel of 56 antibodies with normal human small intestine, colon, and breast tissues. *Tumour Biol.* 19 Suppl 1: 88-99.

Storage Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet Documentation #10041 available at:

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)

HuCAL Anti Mouse IgG2a (HCA037...) [FITC](#), [HRP](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Sheep Anti Mouse IgG (H/L) (AAC10...) [Biotin](#), [FITC](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@800](#), [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Other formats and applications

Gentaur Molecular Products
Voortstraat 49
1910 Kampenhout, Belgium