

***Anti-4-Hydroxy-2-nonenal (4-HNE)
Monoclonal Antibody (HNEJ-2)***

Code: **NNS-MHN-100P-EX**
(100µg of IgG, Lyophilized powder)
NNS-MHN-020P-EX
(20µg of IgG, Lyophilized powder)

Source: **Mouse**

Immunogen: 4-Hydroxy-2-nonenal (4-HNE)-
modified keyhole limpet hemocyanine.

Subclass: IgG1 (κ)

Applications: Immunohistochemistry [see ref.1]
(Recommended concentration: 25 µ g/mL IgG)

Western blotting [see ref.2]
(Recommended concentration: 15 µ g/mL IgG)

Reconstitution: MHN-100P: Dissolve in 1mL of distilled water.
MHN-020P: Dissolve in 1mL of distilled water.


Buffer Concentration: 100 µ g/mL IgG in 50mM Tris-buffered saline (TBS).

Specificity: This antibody show almost negligible reactivity with proteins that were treated with other aldehydes, such as 2-nonenal, 2-hexenal, 1-hexanal, 4-hydroxy-2-hexenal, formaldehyde, or glutaraldehyde. By inhibition test, this antibody has a much higher affinity for the 4-HNE-histidine adduct than 4-HNE-lysine or 4-HNE cysteine adduct.

Storage: Store at less than -20°C.
Avoid repeated freeze & thaw after reconstitution.
For short term storage or transport, storage at 4°C is acceptable.

Stability: 5 years at -20°C.

References: 1) T.Tanaka *et al. Laboratory Investigation* 77(2), p145-155 (1997)
2) S.Toyokuni *et al. FEBS Letters* 359, p189-191 (1995)



For research use only, not for diagnostic use.