



Recombinant Human Interleukin 18 Catalog Number: SJB06

Strength: 25µg

Specifications and Use

Description: Recombinant human IL-18 produced in Yeast is a polypeptide chain containing 157 amino acids, with 4 cys, but no disulfide bond.

Source: Yeast.

Molecular Mass: Approximately 18.3kD.

Purity: ≥97%, as determined by SDS-PAGE and HPLC method.

Endotoxin Level: ≤1EU/µg, determined by the LAL method.

Biological Activity: Induction of IFN-γ by PBMC cell in response to the recombinant human IL-18 was measured using human IFN-γ ELISA. The activity of Lot SJB02-06-1202 was as follow:

IL-18 concentration (ng/ml)	IFN-γ induction (pg/ml)
62.5	30.9
250	40.8

IFN-γ producing activity of the sample cells can be varied depends on cell conditions. Optimal concentration for each application should be determined by each laboratory.

Formulation: Sterile lyophilized powder, with 50mMNaAc, 2mMDTT and 1% sucrose, pH5.5.

Reconstitution: It is recommended that sterile PBS containing at least 0.1% human serum albumin or bovine serum albumin be added to the vial, to prepare a stock solution of not less than 100µg/ml.

Storage: Lyophilized samples are stable for greater than six months from date of receipt at -20° to -70°C.

■ Upon reconstitution, this cytokine can be stored under sterile conditions at 2-8°C for one month or at -20° to -70°C in a manual defrost freezer for three months without detectable loss of activity.

■ **Avoid repeated freeze-thaw cycles.**

Interleukin 18 (IL-18) is a cytokine which identified as a costimulatory factor for production of interferon-γ (IFN-γ) in response to toxic shock and shares functional similarities with IL-12. IL-18 is synthesized as a precursor 24-kDa molecule without a signal peptide and must be cleaved to produce an active molecule. IL-1 converting enzyme (ICE, Caspase-1) cleaves pro-IL-18 at aspartic acid in the P1 position, producing the mature, bioactive peptide that is readily released from the cells. It is reported that IL-18 is produced from Kupffer cells, activated macrophages, keratinocytes, intestinal epithelial cells, osteoblasts, adrenal cortex cells and murine diencephalon. IFN-γ is produced by activated T or NK cells and plays critical roles in the defense against microbial pathogens. IFN-γ activates macrophages and enhances NK activity and B cell maturation, proliferation and Ig secretion. IFN-γ also induces expression of MHC class I and II antigens and inhibits osteoclast activation. IL-18 acts on T helper type-1 (Th1) T cells and in combination with IL-12 strongly induces them to produce IFN-γ. Pleiotropic effects of IL-18 have also been reported, such as, enhancement production of IFN-γ and GM-CSF in peripheral blood mononuclear cells, production of Th1 cytokines, IL-2, GM-CSF and IFN-γ in T cells, enhancement of Fas ligand expression by Th1 cells.

FOR RESEARCH USE ONLY. NOT FOR HUMAN USE.