

## Anti-laminin Monoclonal Antibody

Cat. No.	Quantity	Tested Applications
Y090769	100ul	Immunoprecipitation, Immunoblotting
<b>Antibody Type</b>	Primary Antibody	
<b>Immunogen</b>	PE cells derived from NF-1 teratocarcinoma	
<b>Species Reactivity</b>	mouse	
<b>Specificity</b>	mouse	
<b>Tested Applications</b>	Immunoprecipitation, Immunoblotting	
<b>Application Notes</b>	Optimal dilutions to be determined by researchers	
<b>Raised In</b>	Mouse	
<b>Clonality</b>	Monoclonal	
<b>Isotype</b>	IgG2a	
<b>Purity</b>	Affinity purified	
<b>Storage Buffer</b>	PBS, pH 7.4 with 0.02% sodium azide.	
<b>Formulation</b>	Liquid	
<b>Concentration</b>	10 <sup>10</sup> ug/100ul	
<b>Storage</b>	Store at -20°C/1 year	
<b>References</b>	<p>Wewer, U.M., Tichy, D., Damjanov, A., Paulsson, M., and Damjanov, I. (1987). Distinct antigenic characteristics of murine parietal yolk sac laminin. <i>Dev. Biol.</i> 121, 397-407.</p> <p>Peterson, P.E., Pow, C.S.T., Wilson, D.B., and Hendrickx, A.G. (1993). Distribution of extracellular matrix components during early embryonic development in the Macaque. <i>Acta Anatomica</i> 146, 3-13.</p> <p>Peterson, P.E., Pow, C.S.T., Wilson, D.B., and Hendrickx, A.G. (1993). Characterization of the extracellular matrix during somitogenesis in the long-tailed monkey (<i>Macaca fascicularis</i>). <i>Acta Anatomica</i> 146, 223-233.</p>	

GL